

A new class of models for surface relaxation with exact mean-field solutions

V. Karimipour *

B. H. Seradjeh †

Department of Physics, Sharif University of Technology,
P. O. Box 11365-9161, Tehran, Iran

February 1, 2008

Abstract

We introduce a class of discrete models for surface relaxation. By exactly solving the master equation which governs the microscopic dynamics of the surface, we determine the steady state of the surface and calculate its roughness. We will also map our model to a diffusive system of particles on a ring and reinterpret our results in this new setting.

Keywords: Stochastic process, matrix product ansatz, zero range contact process, surface relaxation.

1 Introduction

Models far from thermodynamic equilibrium, may not obey detailed balance and hence their steady state can not be described by Gibbs Measure, and one often needs to study closely the stochastic dynamics and obtain the steady state directly by simulation, numerical or exact solution of the microscopic dynamical equations. (For recent reviews see [1, 2] and references therein). The microscopic dynamics which is usually encoded in a set of transition rates between different configurations, usually bring about strong long-range correlations in one dimensional systems. This is the case, for example in driven diffusive systems the prototype of which is the Asymmetric Simple Exclusion Process[3]. In such cases then, the mean field solution is not exact or else is exact only under certain conditions on the rates.

*e-mail: vahid@sina.sharif.ac.ir.

†e-mail: babak@sfu.ca

In recent years, much attention has been paid to the exact solution of stochastic processes on one dimensional lattices. Among the methods developed, the Matrix Product Ansatz (MPA), has proved particularly fruitful, since it has both led to simple and nice solutions of the previously known models [4] and has also led to various generalizations [5-12].

The Matrix Product Ansatz is best understood within the operator formalism for Markov processes on one dimensional chains in which the steady state of the process, turns out to be the ground state of a suitable Hamiltonian with nearest neighbor interactions

$$H = \sum_{k=1}^L h_{k,k+1}. \quad (1)$$

According to MPA, the steady state probability $P(\tau_1, \tau_2, \dots \tau_L)$ for the configuration $(\tau_1, \tau_2, \dots \tau_L)$ is written as the trace of a product of noncommuting matrices:

$$P(\tau_1, \tau_2, \dots \tau_L) = \frac{1}{Z(L)} \text{tr} \left(D(\tau_1) D(\tau_2) \dots D(\tau_L) \right) \quad (2)$$

(Here we are taking a closed system with periodic boundary conditions.) It is then easy to show that the above probabilities are the steady state ones, provided that the matrices $D(\tau)$ satisfy a set of relations encoded compactly as:[13]

$$hD \otimes D = X \otimes D - D \otimes X \quad (3)$$

Here D and X are matrix valued column vectors.

The solution of these equations are however far from trivial and so far relatively few systems have lend themselves to this type of solution. In cases where we take $D(\tau)$ to be one dimensional and accordingly denote them by $d(\tau)$, we are in fact searching for exact mean field solutions. Even in this case, (3) is a set of coupled nonlinear algebraic equations and are generally difficult to solve, or else so many constraints of consistency are imposed on the rates to render the process uninteresting and non-physical.

One can however reverse the problem [14, 9, 10, 15] and search among quadratic algebras for those classes that may correspond through the MPA formalism to a stochastic Hamiltonian.

Mathematically, a through search and classification is quite difficult, and one is better to start from a restricted class of algebras or processes which seem to lead to physically interesting models.

(For a partial classification of algebras, which pertain to diffusion and exchange of particles see [15]).

In this paper we take all the matrices to be one dimensional which means we are considering those processes for which mean field solutions are exact. As far as the number of particles of each species on a ring is conserved, a one dimensional representation leads to a rather uninteresting steady state, namely, one in which all the configurations have equal weights. However if there is no such conservation law, then

these mean field solution may be quite nontrivial and contain interesting physics, as we shall see.

Our starting point is to write a set of microscopic dynamical rules pertaining to a generalization of the so-called zero-range contact process [3](see also [16] for more references) and ask if the steady state of the corresponding stochastic process allows an exact mean field solution (i.e: a one dimensional MPA solution). We find that an interesting and rather general class of these processes allow such solutions, namely those processes in which the rates of transitions between neighboring sites can be factorized in two pieces each pertaining to one of the sites. These processes can have two interpretations, one in terms of surface relaxation and the other in terms of diffusive systems of particles. For definiteness we adhere to the first interpretation and determine exactly the steady state of the surface and calculate the roughness of the surface. We study in detail two members of the class.

2 Models for surface relaxation

We consider a ring of L sites, each site of which can accommodate a number of particles on top of each other. At most p particles can be accommodated in each site. To each site k of the lattice ($k = 1, \dots, L$), we assign a stochastic variable τ_k which shows the number of particles in that site, i.e. ($\tau_k = 0, 1, \dots, p$).

The microscopic stochastic dynamics which governs the relaxation of the surface are described according to the following transition rules between the stochastic variables pertaining to any two adjacent sites k , and $k + 1$ (Fig. 1):

$$i \quad j \longrightarrow i - 1 \quad j + 1 \quad \text{with rate} \quad \mu_{i,j}, \quad (4)$$

$$j \quad i \longrightarrow j + 1 \quad i - 1 \quad \text{with rate} \quad \mu_{i,j}, \quad (5)$$

where i and j represent the values of τ_k , and τ_{k+1} . Note that the model has left-right symmetry and we are taking time to be continuous, hence we speak of rates rather than probabilities. Obviously we should have the following constraints on the rates, namely $\mu_{0j} = \mu_{jp} = 0 \quad \forall j$. Moreover we require that the higher the difference between the heights of adjacent columns, the more probable is the above transition. Thus we require that μ_{ij} be an increasing function of $(i - j)$.

Note that the surface does not grow on the average and the total number of particles is constant which we take to be equal to N . If the relaxation of the surface is much faster than the rate of fall of particles, then this model can also describe the growth of the surface, in this case time will be proportional to the number of particles N .

3 Interpretation as a diffusive system of particles

Our model can also be interpreted as a model for particles hopping on a ring. One can now think of L particles on a ring of $L + N$ sites, where each particle k has a number of vacant sites τ_k ahead. In this view, the transition rules (4) are interpreted

as follows: A particle whose distances to its left and (right) neighboring particles are i and (j) , hops forward with rate μ_{ji} and backward with rate μ_{ij}

The increasing property of $\mu_{i,j}$ in terms of $(i - j)$ is still plausible in this setting, specially if we think of traffic flow, since a car in the above condition accelerates if $i - j$ is small and decelerates if $i - j$ is large. In this interpretation the probabilities (2) give the distribution of headways and the roughness designates how uniform the flow is. For reviews on traffic flow and (particularly its statistical mechanical aspects), see [17], and ([18]) respectively.

The above process can now be studied as an interacting stochastic system. For this study we use the operator formalism of Markov processes, where to each configuration $(\tau_1, \tau_2, \dots, \tau_L)$ of the system a state vector $|\tau_1, \tau_2, \dots, \tau_L\rangle$ is assigned and each probability distribution P , is encoded in a vector

$$|P\rangle := \sum_{\tau_1, \tau_2, \dots, \tau_L} P(\tau_1, \tau_2, \dots, \tau_L) |\tau_1, \tau_2, \dots, \tau_L\rangle. \quad (6)$$

The totality of possible state vectors span a complex Hilbert space, denoted by $\mathcal{H} := (C^{p+1})^{\otimes L}$, where C^{p+1} is the complex space spanned by $\mathcal{B} = \{|0\rangle, |1\rangle \dots |p\rangle\}$. The stochastic Hamiltonian describing the above dynamical rules, is given by

$$\mathcal{H} = \sum_{k=1}^L h_{k,k+1}, \quad (7)$$

where

$$h = \sum_{i,j} -\mu_{i,j} \left(|i-1 \ j+1\rangle \langle i \ j| + |j+1 \ i-1\rangle \langle j \ i| - |i \ j\rangle \langle i \ j| - |j \ i\rangle \langle j \ i| \right). \quad (8)$$

Inserting (8) with the choice $\mathbf{D} = \text{column}(d_0, d_1, \dots, d_p)$ and $\mathbf{X} = 0$ into equation (3) we obtain :

$$\mu_{i+1,j-1} d_{i+1} d_{j-1} + \mu_{j+1,i-1} d_{i-1} d_{j+1} = (\mu_{i,j} + \mu_{j,i}) d_i d_j \quad \forall i, j. \quad (9)$$

A sufficient condition for solving these equation is the following:

$$\mu_{j+1,i-1} d_{i-1} d_{j+1} = \mu_{i,j} d_i d_j \quad \forall i, j, \quad (10)$$

For general rates these relations may not have any non-zero solutions, however if we take the rates to be of factorized form, namely

$$\mu_{i,j} = \mu_i \lambda_j, \quad (11)$$

where μ_i and λ_j are arbitrary non-negative real numbers, then the set of equations(10) have the following solution for d_i 's:

$$d_0 = 1, \quad d_1 = \frac{\lambda_0}{\mu_1}, \quad d_2 = \frac{\lambda_1 \lambda_0}{\mu_2 \mu_1} \quad d_3 = \frac{\lambda_2 \lambda_1 \lambda_0}{\mu_3 \mu_2 \mu_1} \quad \dots \quad d_p = \frac{\lambda_{p-1} \dots \lambda_1 \lambda_0}{\mu_p \dots \mu_2 \mu_1} \quad (12)$$

Remark:

The symmetry of the process is crucial in the above derivation, for a totally asymmetric process, the rates will be highly constrained.

Note that the factorization property is the only requirement that exact solvability imposes on this model. We are free to choose the dependence of μ_i and λ_i on i as we wish. It is for physical reasons that we require that

$$\mu_p \geq \mu_{p-1} \cdots \mu_1 \geq \mu_0 = 0 \quad (13)$$

$$\lambda_0 \geq \lambda_1 \geq \cdots \lambda_{p-1} \geq \lambda_p = 0 \quad (14)$$

In this way the rates $\mu_{i,j}$ will be increasing functions of $(i - j)$. Furthermore the probability of a particle jumping from a lower height to a neighboring higher position will be quite low. We can now calculate the probability of all configurations. From (2) we find:

$$P(\tau_1, \tau_2, \cdots \tau_L) = \frac{1}{\mathcal{Z}(L, N)} d(\tau_1) d(\tau_2) \cdots d(\tau_L) \quad (15)$$

Where $\mathcal{Z}(L, N)$, the partition function, is given by:

$$\mathcal{Z}(L, N) := \sum_{\sum_{k=1}^L \tau_k = N} d(\tau_1) d(\tau_2) \cdots d(\tau_L) \quad (16)$$

Due to the commutativity of $d(\tau)$'s, the probabilities depend only on number of columns K_i of a given height i , and not on the position of columns. Thus (15) can be rewritten as:

$$P(K_0, K_1, \cdots, K_p) = \frac{1}{\mathcal{Z}(L, N)} \frac{L!}{K_0! K_1! \cdots K_p!} d_0^{K_0} d_1^{K_1} \cdots d_p^{K_p}, \quad (17)$$

where the combinatorial term account for various possible places of the columns, and (16) can be re-expressed as:

$$\mathcal{Z}(L, N) := \sum'' \frac{L!}{K_0! K_1! \cdots K_p!} d_0^{K_0} d_1^{K_1} \cdots d_p^{K_p} \quad (18)$$

Where by $\sum''$ we mean a sum subject to $\sum_{i=0}^p i K_i = N$, and $\sum_{i=0}^p K_i = L$. Our analysis will be simplified if we consider a grand partition function $\mathcal{Z}_G(L, s)$:

$$\mathcal{Z}_G(L, s) = \sum_{N=0}^{pL} s^N \mathcal{Z}(L, N) \quad (19)$$

$$= \sum_{N=0}^{pL} s^N \sum'' \frac{L!}{K_0! K_1! \cdots K_p!} d_0^{K_0} d_1^{K_1} \cdots d_p^{K_p} \quad (20)$$

$$= (d_0 + d_1 s + \cdots d_p s^p)^L =: (\mathbf{z}(s))^L. \quad (21)$$

The parameter s plays the role of fugacity. The average density of particles $n := \frac{N}{L}$ and the average ratio of columns of height i , that is ; $\overline{k_i} := \frac{\overline{K_i}}{L}$, are given respectively by:

$$n = \frac{s}{L} \frac{\partial}{\partial s} \ln \mathcal{Z}_G(L, s) = s \frac{d}{ds} \ln \mathbf{z}(s) \quad \overline{k_i} = \frac{d_i}{L} \frac{\partial}{\partial d_i} \ln \mathcal{Z}_G(L, s) = \frac{d_i s^i}{\mathbf{z}(s)} \quad (22)$$

The roughness of a surface is usually defined as [19]:

$$\overline{W} := \sqrt{\frac{1}{L} \sum_x \langle (h(x) - \overline{h})^2 \rangle} \quad (23)$$

where $h(x)$ is the height of the surface at point x . This can be rewritten as:

$$\overline{W} = \sqrt{\frac{1}{L} \sum_0^p \overline{K_i} (i - \frac{\overline{N}}{L})^2} \quad (24)$$

A simple calculation shows that:

$$\overline{W}^2 = (s \frac{d}{ds})^2 \ln \mathbf{z}(s) = (s \frac{d}{ds}) n \quad (25)$$

We now consider two examples which lend themselves to elementary analysis.

3.1 Example A

In this first example, we take the rates to be as follows:

$$\mu_i = i \quad \lambda_i = p - i \quad (26)$$

The above assignments satisfy all the requirements put forward on physical grounds. For example the rate of fall of a particle from height p to height 0 is p^2 , while the rate of jump of a particle from a height 1 to $p - 1$ is 1. Moreover $\mu_0 = \lambda_p = 0$. In this case we find from (12):

$$d_i = \frac{p!}{i!(p-i)!} \quad (27)$$

and

$$\mathbf{z}(s) = \sum_{i=0}^p \frac{p!}{i!(p-i)!} s^i = (1+s)^p \quad (28)$$

From (22) we find the average number of particles:

$$n = \frac{ps}{1+s} \quad \text{or} \quad s = \frac{n}{p-n} \quad (29)$$

Also from (22) we find the fraction of sites with height i to be:

$$\overline{k_i} = \binom{p}{i} \frac{s^i}{(1+s)^p} \quad (30)$$

Rewriting this in terms of the number of particles n , we find:

$$\overline{k_i} = \frac{1}{p^p} \binom{p}{i} n^i (p-n)^{p-i}, \quad (31)$$

which implies a binomial distribution for the columns of different heights. The roughness is calculated from (25) and (28) to be:

$$\overline{W} = \sqrt{n(1 - \frac{n}{p})}, \quad (32)$$

If the relaxation is much faster than the rate of fall of particles, we can take $n \sim t$ and $p \sim T$ where T is the time for growing a layer of uniform thickness p , then the above equation implies, that at the early stages of the growth when $n \ll p$, the roughness increases as $\approx t^{\frac{1}{2}}$, in the middle stages of the growth, n is comparable with p , it increases linearly with time, and finally at the final stages when the number of holes $p - n$ becomes a good measure of time $T - t$, the roughness again decreases as the square root of $T - t$.

3.2 Example B:

In this example we consider the following rates:

$$\mu_i = 1 \quad \forall i \neq 0, \quad \lambda_i = 1, \quad \forall i \neq p \quad \mu_0 = \lambda_p = 0 \quad (33)$$

That is, the rates do not depend on the height of source and target columns. We find from (12) that $d_0 = d_1 = \dots d_p = 1$ and thus:

$$\mathbf{z}(s) = \frac{1 - s^{p+1}}{1 - s} \quad (34)$$

from (22) we find:

$$\overline{k_i} = \frac{s^i(1 - s)}{1 - s^{p+1}} \quad (35)$$

Large p -limit Some insight is obtained if we look at the large p limit of the roughness for the two cases $s < 1$ and $s > 1$.

When $s < 1$, we find from (34), that $\mathbf{z}(s) \simeq \frac{1}{1-s}$ and using (22) and (25) we obtain:

$$n \simeq \frac{s}{1-s} \quad \overline{W} \simeq \frac{\sqrt{s}}{(1-s)} = \sqrt{n(1+n)} \quad (36)$$

On the other hand when $s > 1$, we find that $\ln \mathbf{z}(s) \simeq (p+1) \ln s - \ln(s-1)$, which when inserted into (22) and (25), yields respectively

$$n \simeq p - \frac{1}{s-1} \quad \text{and} \quad \overline{W} \simeq \frac{\sqrt{s}}{s-1} = \sqrt{(p-n)(p-n+1)} \quad (37)$$

Moreover from (35) we find that in the large p limit, when $s < 1$ or $n < p/2$,

$$\overline{k_i} \simeq \frac{n^i}{(1+n)^{i+1}} \quad (38)$$

and when $s > 1$ or $n > p/2$

$$\overline{k_i} \simeq \frac{(p-n)^{p-i}}{(1+p-n)^{p-i+1}} \quad (39)$$

It is interesting to note that despite the very different nature of the transition rates in models **A** and **B**, in both models, the formulas obtained for the roughness imply that the roughness first increases as $t^{1/2}$ and then increases linearly and finally decreases at the final stage again as $(T-t)^{1/2}$.

These two regimes are separated by the case when $s = 1$, or $n = \frac{p}{2}$, where we find for all arbitrary p 's, $\overline{k_i} = \frac{1}{p+1}$, meaning that all heights are equally probable. In this case the roughness turns out from (25) to be:

$$\overline{W} = \sqrt{\frac{p^2}{12} + \frac{p}{6}}. \quad (40)$$

Fig.(3) shows the value of roughness in terms of the fugacity for $p = 10$. It is seen that there is a sharp peak in the roughness at $n = \frac{p}{2}$, when half of the capacity of the lattice is filled with particles.

All these hint that the point $s = 1$ or $n = \frac{p}{2}$ is a special point, possibly a point of phase transition.

Acknowledgement: V. K. would like to thank V. Rittenberg for his valuable comments. We would like to thank A. Rezakhani for his valuable help in preparing the manuscript.

References

1. H. Hinrichsen: Critical phenomena in non-equilibrium systems, cond-mat/0001070.
2. G. M. Schütz, in *Phase Transitions and Critical Phenomena* edited by C. Domb and J. Lebowitz (Academic Press,London, 1999).
3. F. Spitzer, Adv. Math. **5**,246(1970).
4. B. Derrida, M.R. Evans, V.Hakim and V. Pasquier, J. Phys. A: Math. Gen. **26**,1493, (1993).
5. P. Arndt, T. Heinzel and V. Rittenberg; J. Phys. A; Math. Gen. **31**, 833,(1998).
6. M. R. Evans; J. Phys. A: Math. Gen.**30**, 5669 (1997); Europhys. Lett.**36**,13, (1996).
7. M. R. Evans, D. P. Foster, C. Godreche D. Mukamel; J. Stat. Phys. **80** (1995); Phys. Rev. Lett. **74**, 208 (1995).
8. H.W. Lee, v. Popkov, and D. Kim, J. Phys. A **30**, 8497 (1997).
9. V. Karimipour, Phys. Rev. E **59**205 (1999).
10. V. Karimipour, Europhys. Letts. **47**(3), 304(1999).
11. M. Khorrami, V. Karimipour; J. Stat. Phys. **100** 5/6,(2000) 999.
12. J. de Gier and B. Nienhuis, Phys. Rev. E **59**,4899,(1999).
13. K. Krebs and S. Sandow; J.Phys. A ; Math. Gen. **30** 3165(1997).
14. F. C. Alcaraz, S. Dasmahapatra and V. Rittenberg V J.Phys. A ; Math. Gen. **31** 845 (1998).
15. A. P. Isaev, P. N. Pyatov, and V. Rittenberg, Diffusion algebras, to be published.
16. M. R. Evans; Phase transitions in one-dimensional non-equilibrium systems, cond-mat/0007293.
17. D. Helbing, Traffic and related self-driven many particle systems, cond-mat/0012229.
18. D. Chowdhury, L. Santen, and A. Schadschneider, *Phys. Rep.***329**,199 (2000).
19. A. L. Barabasi, and H. E. Stanley; *Fractal Concepts in Surface Growth*Cambridge University Press, (1994).




Figure 1: In the surface relaxation model, a particle drops with rate μ_{ij}




Figure 2: In the diffusive system, a particle hops with rate μ_{ij}




Figure 3: The square of roughness for $p=10$